



Estimation of the magnetic field gradient tensor using the Swarm constellation

Kotsiaros, Stavros; Finlay, Chris; Olsen, Nils

Published in:
Geophysical Research Abstracts

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Kotsiaros, S., Finlay, C., & Olsen, N. (2014). Estimation of the magnetic field gradient tensor using the Swarm constellation. *Geophysical Research Abstracts*, 16, [EGU2014-5611].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Estimation of the magnetic field gradient tensor using the Swarm constellation

Stavros Kotsiaros, Chris Finlay, and Nils Olsen
DTU Space, Denmark (skotsiaros@space.dtu.dk)

For the first time, part of the magnetic field gradient tensor is estimated in space by the Swarm mission. We investigate the possibility of a more complete estimation of the gradient tensor exploiting the Swarm constellation. The East-West gradients can be approximated by observations from the lower pair of Swarm satellites, whereas the North-South gradients can be approximated by the first differences in the along-track direction. As a preliminary test, here we will present an analysis of along track differences of CHAMP vector observations. These show considerably smaller standard deviations compared to conventional vector observations at almost all latitudes. Analytical and numerical analysis of the spectral properties of the gradient tensor shows that specific combinations of the East-West and North-South gradients have almost identical signal content to the radial gradient. Gradient data are less contaminated by large scale fields produced in the magnetosphere and ionosphere and specific gradient combinations can lead to an improved determination of both the lithospheric field and the high degree secular variation.